

# Production protocol Furosemide suspension (10 mg/mL) with Liquifine® Alkalic Dry



## Preparation instructions

1. Weigh the required quantity of Liquifine® Alkalic dry (6.3 g per 100 ml suspension) and transfer it to the final recipient<sup>(\*)</sup>.
  2. Weigh the required quantity of furosemide and transfer it also to the final recipient.
  3. Shake firmly.
  4. Add purified water to the desired volume and shake firmly until a homogeneous dispersion is obtained.
- (\*) You can also use the pre-filled 100 mL bottles with mark line, which contain already the required amount of Liquifine® Alkalic dry.

## Production report

Preparation date	/ /	Galenic type	Suspension
Batch number		Volume	
Raw material	Number analysis register	Required Quantity	Weighed quantity
Furosemide		g	g
Liquifine® Alkalic dry		g	g
Purified water		Ad mL	Ad mL

## Tarra

• \_\_\_\_\_

## Quality control

• Homogeneity: \_\_\_\_\_

## Stability

• 60 days

## Storage and labelling

- Protected from light – Fridge
- Shake before use

Name and Signature Compounder	Name and Signature Pharmacist

## MAGIS PHARMA

Keizershoek 336, 2550 Kontich, T: +32 3 457 11 76, F: +32 3 457 43 40  
info@magis-pharma.be, www.magis-pharma.be